

URINARY S/O

LOWER URINARY TRACT DISEASE



URINARY S/O



LOAF

Available in 200 g, 410 g can

RECOMMENDED DAILY INTAKE & FEEDING DURATION

DOG'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	can	grams	can	grams	can
2	208	2/4	183	2/4	158	2/4
4	350	3/4	308	3/4	266	3/4
6	475	1 + 1/4	418	1	361	1
8	589	1 + 2/4	519	1 + 1/4	448	1
10	697	1 + 3/4	613	1 + 2/4	529	1 + 1/4
15	944	2 + 1/4	831	2	718	1 + 3/4
20	1172	1 + 3/4	1031	2 + 2/4	890	2 + 1/4
25	1385	3 + 2/4	1219	3	1053	2 + 2/4
30	1588	3 + 3/4	1397	3 + 2/4	1207	3
35	1783	4 + 1/4	1569	3 + 3/4	1355	3 + 1/4
40	1970	4 + 3/4	1734	4 + 1/4	1498	3 + 3/4
45	2152	5 + 1/4	1894	4 + 2/4	1636	4
50	2329	5 + 3/4	2050	5	1770	4 + 1/4
55	2502	6	2202	5 + 1/4	1902	4 + 3/4
60	2671	6 + 2/4	2350	5 + 3/4	2030	5
70	2998	7 + 1/4	2638	6 + 2/4	2279	5 + 2/4
80	3314	8	2916	7	2519	6 + 1/4

ADULT DOG

Urinary S/O is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

RECOMMENDATIONS: It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

NOTE: in older dogs, the use of Urinary S/O Ageing* is recommended and renal function should be checked before the advise of a new diet. It is recommended to divide the daily intake into 2 meals.

RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



LOW RSS

Helps lower the concentration of ions contributing to crystal formation.

[S/O® INDEX] This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	83.68 g	8 g
Fat content	62.76 g	6 g
Carbohydrate	48.12 g	4.6 g
Crude fibre	14.64 g	1.4 g
Dietary fibre	25.1 g	2.4 g
Calcium	2.2 g	0.21 g
Phosphorus	2.09 g	0.2 g
Potassium	2.09 g	0.2 g
Sodium	3.03 g	0.29 g
Chloride	4.18 g	0.4 g
Magnesium	0.17 g	0.02 g
Copper	3.14 mg	0.3 mg
Zinc	23.01 mg	2.2 mg
Taurine	1.26 g	0.12 g
Vitamin A	47071.13 IU	4500 IU
Vitamin E	198.74 mg	19 mg
Omega 6	25.1 g	2.4 g
Omega 3	1.36 g	0.13 g
EPA + DHA	0.63 g	0.06 g
Metabolisable energy*	96	

*Calculated according to NRC2006.

COMPOSITION

Meat and animal derivatives (chicken and pork meat), oils and fats (sunflower oil, fish oil), cereals (corn flour, corn starch), derivatives of vegetable origin (cellulose, gelling agent, marigold extract), minerals, various sugars (caramel). No animal medicine included.

ADDITIVES (KG)**

Nutritional additives: Vitamin D3: 190 IU, E1 (Iron): 3.4 mg, E2 (Iodine): 0.24 mg, E4 (Copper): 2 mg, E5 (Manganese): 1 mg, E6 (Zinc): 10 mg, Sulfur: 0.2%, Urine acidifying substances: calcium sulfate: 0.25%. Technological additives: Sodium bisulfate: 0.56 g.

**Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

