

URINARY S/O

SMALL DOGS



Available in 1.5 kg, 4 kg, 8 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

DOG'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	cups	grams	cups	grams	cups
2	55	5/8	48	4/8	42	4/8
2,5	65	5/8	57	5/8	49	4/8
3	74	6/8	65	5/8	56	5/8
3,5	83	7/8	73	6/8	63	5/8
4	92	1	81	7/8	70	6/8
5	109	1 + 1/8	96	1	83	7/8
6	125	1 + 2/8	110	1 + 1/8	95	1
7	140	1 + 4/8	123	1 + 2/8	106	1 + 1/8
8	155	1 + 5/8	136	1 + 3/8	117	1 + 2/8
9	169	1 + 6/8	149	1 + 4/8	128	1 + 3/8
10	183	1 + 7/8	161	1 + 5/8	139	1 + 4/8

Urinary S/O Small Dog is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

RECOMMENDATIONS: It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Small Dog for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

NOTE: in older dogs, the use of Urinary S/O Ageing* is recommended and renal function should be checked before the advise of a new diet. It is recommended to divide the daily intake into 2 meals.

LOWER URINARY TRACT DISEASE



SMALL DOGS

RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



TARTAR CONTROL

Nutrient that traps the calcium in saliva so reducing tartar deposits.

S/O® INDEX This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	51.98 g	20 g
Fat content	44.18 g	17 g
Carbohydrate	114.09 g	43.9 g
Crude fibre	5.98 g	2.3 g
Dietary fibre	14.55 g	5.6 g
Calcium	1.85 g	0.71 g
Phosphorus	1.82 g	0.7 g
Potassium	2.08 g	0.8 g
Sodium	3.12 g	1.2 g
Chloride	5.2 g	2 g
Magnesium	0.13 g	0.05 g
Copper	3.9 mg	1.5 mg
Zinc	44.18 mg	17 mg
Taurine	0.49 g	0.19 g
Vitamin A	4547.82 IU	1750 IU
Vitamin E	124.74 mg	48 mg
Omega 6	9.3 g	3.58 g
Omega 3	1.72 g	0.66 g
EPA + DHA	0.57 g	0.22 g
Metabolisable energy*	385	

*Calculated according to NRC2006.

COMPOSITION

Rice, maize flour, dehydrated poultry protein, animal fats, maize gluten, hydrolysed animal proteins, minerals, vegetable fibres, egg powder, wheat gluten**, soya oil, fish oil, fructo-oligo-saccharides, marigold extract (source of lutein). Protein sources : dehydrated poultry protein, maize gluten, hydrolysed animal proteins, egg powder. **L.I.P.: protein selected for its very high digestibility.

ADDITIVES (KG)***

Nutritional additives: Vitamin A: 16000 IU, Vitamin D3: 1000 IU, E1 (Iron): 47 mg, E2 (Iodine): 4.7 mg, E4 (Copper): 14 mg, E5 (Manganese): 61 mg, E6 (Zinc): 148 mg, E8 (Selenium): 0.06 mg, Sulphur: 0.5%, Urine acidifying substance: calcium sulphate (0.8%). Technological additives: Pentasodium triphosphate: 3.5 g. Preservatives - Antioxidants.

***Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O

